MOORED OBSERVATIONS FOR THE FORCED UPPER OCEAN DYNAMICS EXPERIMENT IN THE ARABIAN SEA

John Marra
Lamont-Doherty Earth Observatory,
PO Box 1000,
Palisades, NY 10964
Tel: 941-365-8891, Fax: 914-365-8150,
e-mail, marra@ldeo.columbia.edu

Award # N00014-94-1-0450

LONG-TERM GOALS

My long-term goals are to understand how biological processes in the ocean affect, and are affected by, physical processes (mixing, stirring, and the penetration of light into the ocean). I want to figure out how irradiance and wind-stress influence the dynamics of the upper ocean, how these dynamics govern the rate and nature of primary production, and ultimately, the growth of phytoplankton populations. I also want to understand the feedback between the changes in phytoplankton biomass and the variability in optical properties of the sea, and to use the absorption and scattering properties of phytoplankton to predict photosynthesis and growth.

OBJECTIVES

The scientific objectives of this grant are to (1) determine the diurnal, within-season, and seasonal variability of phytoplankton for a locale subject to strong seasonal wind-stress; (2) understand how POC:Chl changes with the advent of the SW monsoon; (3) determine whether the seasonal increase in phytoplankton affects the surface heating of the water column; and (4) determine whether the chlorophyll specific primary production is constant through the year in the Arabian Sea, where strong seasonal signals in nutrients are characteristic.

WORK COMPLETED

- 1. We completed two data reports, one for each deployment (Ho et al., 1996a, 1996b), and contributed to a preliminary results article for EOS (Rudnick et al., 1997).
- 2. We made presentations of data at the Arabian Sea Workshop in July, 1997.
- 3. We submitted two manuscripts for the Deep-Sea Research-II Special Volume on the Arabian Sea Expedition (Marra et al., submitted; Kinkade et al., submitted) and were coauthors on three other manuscripts.
- 4. Abstracts were submitted for the Ocean Sciences Meeting, held in February, 1998.

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send comment arters Services, Directorate for Inf	s regarding this burden estimate formation Operations and Reports	or any other aspect of the property of the contract of the con	his collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 30 SEP 1997	2 DEDORT TYPE			3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
Moored Observations for the Forced Upper Ocean Dynamics Experiment in the Arabian Sea				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Lamont-Doherty Earth Observatory of Columbia University, Palisades, NY, 10964				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAII Approved for publ	ABILITY STATEMENT ic release; distributi	ion unlimited			
13. SUPPLEMENTARY NO	TES				
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFIC		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	4	

Report Documentation Page

Form Approved OMB No. 0704-0188

RESULTS

The year-long mooring records show clearly the semi-annual biological response to the northeast and southwest monsoons. Interestingly, prior to each mixed-layer deepening associated with the monsoons, there is evidence of mesoscale eddies passing the mooring site, and which contain enhanced quantities of chlorophyll. Each monsoon causes the mixed layer to deepen, but for the northeast monsoon, convective processes dominate, while during the southwest monsoon, the mixed layer deepens via wind-induced mixing. During both mixing periods, chlorophyll declines, and then increases as the water column stratifies. The annual cycle of chlorophyll variability at the mooring site (the climatological center of the Findlater Jet, 15°30'N/61°30'E) generally follows the mixed layer depth. When the mixed layer is deep, the quantity of chlorophyll in the water column is low, and increases when the mixed layer shallows following the cessation of monsoon winds. Prolonged stratification, during the intermonsoons, also results in declines in photosynthesis.

The variability in productivity over the year generally follows the change in chlorophyll (see Fig. 1 from Marra et al., 1997), except where there may have been changes to community structure (Feb-Mar, '95), or when grazing might be important (May, July, '95).

Strong diel variability in phytoplankton and particle variables were observed, and coherence analysis indicated that these were modulated more by water column processes (e.g., changes in the mixed layer depth) than by solar variability (Kinkade et al., 1997).

IMPACT

We have demonstrated the utility of moored sensor systems in an unforgiving environment. The high-resolution data are extremely useful to those trying to understand the drivers for seasonal variability in productivity. The seasonal variability in productivity is crucial to interpret the export of carbon from the surface ocean to depth. The results will fuel consideration of the importance of diel processes in determining the activities of plankton. We now have a better idea of the relationships among phytoplankton production, nutrient supply, and changes in mixed layer depth. The importance of mesoscale processes in distributing heat and momentum was demonstrated over a large part of the study area, and is probably the reason why productivity can remain relatively high despite the extreme variability in wind-forcing.

TRANSITIONS

During this project we have developed a low-cost, compact, easy-to-use moored spectral-radiometer with 32 wavelengths. It should find a variety of uses. The most recent issue of Sea Technology (Sept., '97) has advertisements by four companies selling *in situ* fluorometers. I think that the efforts by both the LDEO and UCSB (T. Dickey et al.) have

had something to do with the use of chlorophyll fluorescence measurements at sea. We have also continued to improve our bio-optical models for calculations of primary production, which enhances predictability of optical variability.

RELATED PROJECTS

We are collaborating with others in the overall Arabian Sea Program (including JGOFS), contributing to studies of shipboard productivity measurements (R.T. Barber and myself), the moored sediment trap program (e.g., C. Lee and S. Honjo), air-sea interaction (R. Weller, A. Fischer). I am continuing comparisons of results with previous mooring programs, primarily to examine further diel variability in the moored-sensor variables with other kinds of rate measurements.

REFERENCES

Ho, C., C. S. Kinkade, C. Langdon, M. Maccio and J. Marra. 1996a. The Forced Upper Ocean Dynamics Experiment in the Arabian Sea. Results from the multi-variable moored sensors from deployment-2 of the WHOI mooring. LDEO Tech Report LDEO-96-7, pp. 19+figs., app.

Ho, C., C.S. Kinkade, C. Langdon, M. Maccio and J. Marra. 1996b. The Forced Upper Ocean Dynamics Experiment in the Arabian Sea. Results from the multi-variable moored sensors from deployment-2 of the WHOI mooring. LDEO Tech Report LDEO-96-8, pp. 19+figs., app.

Kinkade, C.S., J. Marra, T. Dickey, C. Langdon, D. Sigurdson, and R. Weller. 1997. Diel variability of bio-optical signals in the Arabian Sea. Deep-Sea Research-II (submitted).

Marra, J., T.D. Dickey, C. Ho, C.S. Kinkade, D.E. Sigurdson, R. Weller and R. T. Barber. 1997. Variability in primary production as observed from moored observations in the central Arabian Sea in 1995. Deep-Sea Research-II (submitted).

Rudnick, D., R.A. Weller, C. Eriksen, T.D. Dickey, J. Marra and C. Langdon. 1997. One-year moored observations of the Arabian Sea Monsoons. EOS 78, pp. 117, 120-121.




Figure 1. Areal photosynthetic carbon assimilation calculated from a bio-optical model plotted against areal chlorophyll a determined from the moored fluorescence sensors (and other data), from Marra et al. (1997).